



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$(x - a_1)^n + \binom{n}{2}(a_2 - a_1^2)x^{n-2} + \varphi(x) = 0,$$

where $\varphi(x)$ is a polynomial of degree $n - 3$ at most. Suppose $a_2 = a_1^2$, and write $y = x - a_1$. The equation becomes

$$y^n + \varphi(y + a_1) \equiv y^n + c_3y^{n-3} + \dots + c_n = 0.$$

If all the c 's are zero, the theorem is true. If not, let c_k be the first coefficient not zero, and c_e the last. The equation is then

$$y^n + c_ky^{n-k} + \dots + c_e y^{n-l} = 0.$$

Apply Descartes's rule to this equation. If we call the left hand side $f(y)$, $f(y)$ and $f(-y)$ can have together at most $n - k + 1 - l$ variations of sign (k odd), or $n - k + 2 - l$ (k even), hence at most $n - 2 - l$. The equation has, therefore, at least $2 + l$ zero or complex roots. Exactly l roots are zero, however, hence it must have 2 complex. This contradicts our hypothesis that the roots were all real. Hence all the c 's are zero and $f(y) \equiv y^n$, and the equation in x is $(x - a_1)^n = 0$.

Also solved by LAENAS G. WELD.


GEOMETRY.

456. Proposed by J. W. CLAWSON, Ursinus College.

The interior and exterior bisectors of the angles A, B, C of a triangle meet the opposite sides in $U, U'; V, V'; W, W'$ respectively. Circles are drawn on UU', VV', WW' as diameters (Circles of Apollonius). Prove that (1) these three circles have a common chord. (2) The center of the circumcircle lies on this common chord.

SOLUTION BY PROPOSER.

1. $A(BC, UU')$ is a harmonic pencil; so (BC, UU') is a harmonic range. If T be any point on the circle having UU' as diameter, $T(BC, UU')$ is a harmonic



pencil. But UTU' is a right angle. Therefore TU bisects $\angle BTC$. Hence,

$$BT : TC = BU : UC = BA : AC.$$

In particular, if the circles having UU' and WW' as diameters intersect at the points P and Q ,

$$BP : PC = BA : AC \text{ and } BQ : QC = BA : AC.$$

Similarly,

$$AP : PB = AC : CB \text{ and } AQ : QB = AC : CB.$$

Hence,

$$AP : PC = AB : BC \text{ and } AQ : QC = AB : BC.$$

It follows that P and Q are points on the circle having VV' as diameter. Hence the three circles UU' , VV' , WW' are coaxial.

2. Since (BC, UU') is a harmonic range, any circle passing through B and C is orthogonal to the circle having UU' as diameter. Hence, the circumcircle is orthogonal to that circle. Similarly the circumcircle is orthogonal to the circles VV' and WW' .

Hence, if O be the circumcenter, OA is a tangent to UU' , OB to VV' and OC to WW' . But these tangents are equal in length.

Hence O is a point on the common chord of the circles UU' , VV' , WW' .

Also solved by C. E. HORNE, DAVID F. KELLEY, ELIJAH SWIFT, ROGER A. JOHNSON, and C. N. SCHMALL.


457. Proposed by NATHAN ALTSHILLER, University of Washington.

AB and AC are respectively a diameter and a chord of a circle whose center is O . The lines joining B to the extremities of the diameter perpendicular to AC , meet AC in the points M, N . Express the angle MON in terms of the angle CAB .

SOLUTION BY J. A. CAPARO, University of Notre Dame.

Let E and D be the extremities of the diameter perpendicular to AC at the point K . With the notation of the figure we easily see that

$$\sin \angle AMB = \sin \angle ABN = \cos z.$$



Hence, by the sine law, letting $AB = 2R$, we have in $\triangle ANB$, $AN = 2R \cot z$, and in $\triangle AMB$, $AM = 2R \tan z$. Also from $\triangle AKO$, $AK = R \cos y$, $OK = R \sin y$. $KN = AN - AK = 2R \cot z - R \cos y$, $MK = AM - AK = 2R \tan z - R \cos y$.

$$\therefore \tan u = \frac{MK}{OK} \frac{2 \tan z - \cos y}{\sin y}. \quad \text{Also } \tan (x + u) = \frac{KN}{KO}.$$